

- (c) For the linear transformation $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined as, $L(v) = A \cdot v$, where

$$A = \begin{pmatrix} -9 & 2 & 1 \\ -6 & 1 & 1 \\ 5 & 0 & -2 \end{pmatrix}$$

Determine, whether L is an isomorphism or not.
(7.5)

[This question paper contains 8 printed pages.]

Your Roll No.....

Sr. No. of Question Paper : 4945

H

Unique Paper Code : 2352571201

Name of the Paper : Elementary Linear Algebra /

Name of the Course : **B.A./B.Sc. (Prog.) with
Mathematics as Non-Major/
Minor – DSC**

Semester : II

Duration : 3 Hours

Maximum Marks : 90

Instructions for Candidates

1. Write your Roll No. on the top immediately on receipt of this question paper.
2. Attempt **all** questions by selecting **any two** parts from each question.
3. **All** questions carry equal marks.
4. Use of simple calculator is allowed.

4945

2

1. (a) If x and y are vectors in $\mathbb{R}^n$, then prove that

$$|x \cdot y| \leq \|x\| \|y\|. \text{ Also verify it for the vectors}$$

$$x = [2, 2, 1] \text{ and } y = [-4, 0, 3] \text{ in } \mathbb{R}^3. \quad (5.5+2)$$

- (b) Solve the following system of linear equations using

Gaussian Elimination method. Also indicate whether

the system is consistent or inconsistent.

$$-5x - 2y + 2z = 14$$

$$3x + y - z = -8$$

$$2x + 2y - z = -3 \quad (6.5+1)$$

- (c) Find the quadratic equation $y = ax^2 + bx + c$

that goes through the points $(-2, 20)$, $(1, 5)$ and

$$(3, 25). \quad (7.5)$$

4945

7

- (c) Define a linear transformation from a vector space

V to W . If $T : V \rightarrow W$ is a linear transformation

and S is a subspace of W , then show that the set

$$T^{-1}(S) = \{v \in V \mid T(v) \in S\} \text{ is a subspace of } V.$$

$$(2+5.5)$$

6. (a) Consider the linear transformation $L : M_{22} \rightarrow M_{32}$ defined as :

$$L\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}\right) = \begin{bmatrix} a & 0 \\ 0 & 0 \\ 0 & b \end{bmatrix}, \text{ show that } \dim(\text{Range}(L)) +$$

$$\dim(\text{Ker}(L)) = \dim(M_{22}). \quad (7.5)$$

- (b) Let $L : P(x) \rightarrow P(x)$ be a linear operator defined

as $L(p(x)) = x p(x)$, where $P(x)$ denotes the vector

space of real polynomials. Show that L is one-one

but not onto. $(4+3.5)$

P.T.O.

5. (a) Check if the following mappings defined on M_{22} the set of 2×2 real matrices, are linear transformation or not. Prove it or give a counter example to disprove.

(i) $f: M_{22} \rightarrow \mathbb{R}$ defined as $f(A) = \text{trace}(A)$,
where $\text{trace}(A)$ is the sum of diagonal elements of the matrix A .

(ii) $g: M_{22} \rightarrow \mathbb{R}$ defined as $g(A) = \det(A)$.
(4+3.5)

- (b) Find the matrix of linear transformation $f: P_3(x) \rightarrow \mathbb{R}^3$ with respect to standard ordered bases defined as:

$$f(ax^3 + bx^2 + cx + d) = (a + 2b - c, 2b + d, a - c + d)$$

(7.5)

2. (a) Find the reduced row echelon form of the following matrix : (7.5)

$$\begin{bmatrix} 1 & -3 & 2 & -4 & 8 \\ 3 & -9 & 6 & -12 & 24 \\ -2 & 6 & -5 & 11 & -18 \end{bmatrix}$$

- (b) Determine whether the vector $[7, 1, 18]$ is in the row space of the given matrix:

$$\begin{bmatrix} 3 & 6 & 2 \\ 2 & 10 & -4 \\ 2 & -1 & 4 \end{bmatrix} \quad (7.5)$$

- (c) Find the characteristic polynomial, eigenvalues and corresponding eigenvectors for the given matrix:

$$\begin{bmatrix} 2 & -3 \\ 0 & 2 \end{bmatrix} \quad (2+2+3.5)$$

3. (a) Use the Diagonalization Method to determine whether the following matrix is diagonalizable.

$$\begin{bmatrix} 7 & 1 & -1 \\ -11 & -3 & 2 \\ 18 & 2 & -4 \end{bmatrix} \quad (7.5)$$

- (b) Show that the set P_3 of all real polynomials of degree ≤ 3 is a vector space under the usual (term-by-term) operations of addition and scalar multiplication. (7.5)

- (c) Define span of S , where S a nonempty subset of a vector space V . Determine span (S) where $S = \{(1,1,0), (2,1,3)\}$ is a subset of R^3 . Also examine whether the following vectors of R^3 are in span (S): (i) $(0,0,0)$; (ii) $(1,2,3)$. (2+3.5+1+1)

4. (a) Define a linearly independent set of vectors in a vector space. Check whether the following subsets of R^3 are linearly independent or not.

(i) $\{(1,2,3), (2,3,1), (3,5,4)\}$

(ii) $\{(1,0,0), (0,0,-5)\}$ (2.5+2.5+2.5)

- (b) Define a finite dimensional vector space. Let W be the solution set to the matrix equation $AX = 0$,

where $A = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \end{pmatrix}$.

Show that (i) W is a subspace of R^3 .

(ii) Find a basis for W . (2+3+2.5)

- (c) Let P_2 be the vector space of all real polynomials of degree ≤ 2 . Show that $S = \{1, x + 1, 2x + x^2\}$ is a basis of P_2 . (7.5)