

[This question paper contains 12 printed pages.]

Your Roll No.....

Sr. No. of Question Paper : 2122

I

Unique Paper Code : 2922071103

Name of the Paper : Statistics for Business Decisions

Name of the Course : **Bachelor of Business Administration (Financial Investment Analysis)**

Semester : I

Duration : 3 Hours

Maximum Marks : 90

Instructions for Candidates

1. Write your Roll No. on the top immediately on receipt of this question paper.
2. Answer any **five** questions.
3. **All** Questions carry equal marks which are indicated on the side in bold numbers.
4. Use of Simple Calculator is allowed.
5. Reference to statistics tables is allowed.

1. (a) State true or false :

- (i) The complete collection of all entities under study is called the sample.
- (ii) A census is the process of gathering data on all the entities in the population.
- (iii) A variable is a numerical description of each of the possible outcomes of an experiment.
- (iv) The lowest level of data measurement is the ratio level.
- (v) The main objective of business analytics is to transform data into meaningful information for business managers.

P.T.O.

- (vi) An event that cannot be broken down into other events is called a certainty outcome.
- (vii) A joint probability is the probability that at least one of two events occurs.
- (viii) Bayes' rule is a rule to assign probabilities under the relative frequency method.
- (ix) If the mean, median, and mode are equal, then the distribution is positively skewed. $(1 \times 9 = 9)$
- (b) The table given below presents data regarding sales in thousands and advertisement in million rupees for the last fifteen years.

Year	Sales (000 units)	Advertisement (Rs. million)
2012	187	49
2013	186	45
2014	200	56
2015	191	52
2016	192	58
2017	192	47
2018	193	50
2019	193	52
2020	215	52
2021	201	59

Find out mean, median, variance and standard deviation for sales and advertisement. (9)

2. (a) Consumer Reports tested 166 different point-and-shoot digital cameras. Based upon factors such as the number of megapixels, weight (oz.), image quality, and ease of use, they developed an overall score for each camera tested. The overall score ranges from 0 to 100, with higher scores indicating better overall test results. Selecting a camera with many options can be a difficult process, and price is certainly a key issue for most consumers. By spending more, will a consumer really get a superior camera? And, do cameras that have more megapixels, a factor often considered to be a good measure of picture quality, cost more than cameras with fewer megapixels? Table given below shows the brand, average retail price (\$), number of megapixels, weight (oz.), and the overall score for 13 Canon and 15 Nikon subcompact cameras tested by Consumer Reports.

Obs.	Brand	Price (\$)	Megapixels	Weight (oz.)	Score
1	Canon	330	10	7	60
2	Canon	200	12	5	66
3	Nikon	300	12	7	60
4	Canon	200	10	6	62
5	Canon	180	12	5	62
6	Canon	200	12	7	61
7	Canon	200	14	5	60
8	Nikon	130	10	7	60
9	Nikon	130	12	5	59
10	Nikon	130	16	5	50

Using simple linear regression, develop an estimated regression equation that could be used to predict the price given the score of the camera. Also compute coefficient of determination. Estimate price of a camera with a score of 80. (9)

- (b) The amount of loan granted per borrower in a small co-operative bank is normally distributed with a mean of Rs. 28,000 and standard deviation of Rs. 6,400. If a bank borrower is randomly selected, find the probability that the loan amount is
- (i) More than Rs. 22,000;
 - (ii) Less than Rs. 10,000;
 - (iii) Between Rs. 21,000 and Rs. 30,000.
 - (iv) The bank decided to give some relaxation in rate of interest to 4% of the smallest borrowers. What should be the loan amount to be eligible for this relaxation?
 - (v) 25% of all the loans are less than what amount? (9)
3. (a) Ship collisions in the Houston Ship Channel are rare. Suppose the number of collisions are Poisson distributed, with a mean of 1.2 collisions every four months. What is the probability of having one or fewer collisions in a six-month period? [Given $\exp(-1.2) = 0.301194$; $\exp(-1.5) = 0.22313$; $\exp(-1.8) = 0.165299$] (9)
- (b) Delhi has been experiencing poor air quality in winters since last five years. An Air Quality Index (AQI) more than 300 is considered 'very poor'. Various newspapers are reporting that AQI is more than 300 in Delhi this winter. However, the Government is claiming that because of the various measures taken by it, the AQI is less than 300. To verify its claim, the government measures AQI from 40 sample places and finds that AQI is 297. If σ of AQI is 18, is the government's claim valid at 5% level of significance? Also compute p-value to conclude your answer. (9)

4. (a) Differentiate between correlation and Regression analysis. If standard deviation of X and Y are 4 and 3 respectively, and coefficient of correlation between X and Y is 0.8, find the two regression coefficients ('Y on X' and 'X on Y'). (9)

- (b) The editor-in-chief of a major metropolitan newspaper has been trying to convince the paper's owner to improve the working conditions in the pressroom. He is convinced that the noise level when the presses are running creates unhealthy levels of tension and anxiety. He recently had a psychologist conduct a test during which press operators were placed in rooms with varying levels of noise and then given a test to measure mood and anxiety levels. The following table shows the index of their degree of arousal or nervousness and the level of noise to which they were exposed (1.0 is low and 10.0 is high).

Noise level	4	3	1	2	6	7	2	3
Degree of arousal	39	38	16	18	41	45	25	38

- (i) Develop an estimating equation that describes these data.

- (ii) Predict the degree of arousal we might expect when the noise level is 5. (9)

5. (a) Department of Financial Studies is planning a trip in the last week of October. Weather forecast indicates that there is a possibility of rain/hailstorm/thundering in October and chances of each are 20%, 35% and 45%.

Past data suggests that on 50% of the occasions whenever there was a rain, the trip got cancelled, on 60% of times when there was a hailstorm the trip got cancelled and 40% of the times whenever there was thundering the trip got cancelled.

The recently planned trip just got cancelled. What is the probability that it cancelled due to rain? (9)

- (b) Market researcher Gartner Dataquest reports that Dell Computer controls 27% of the PC market in Delhi. Suppose a business researcher randomly selects 130 recent purchasers of PC.

- (i) What is the probability that more than 39 PC purchasers bought a Dell computer?
- (ii) What is the probability that between 28 and 38 PC purchasers (inclusive) bought a Dell computer?
- (iii) What is the probability that fewer than 23 PC purchasers bought a Dell computer?
- (iv) What is the probability that exactly 33 PC purchasers bought a Dell computer? (9)

6. Attempt **any two** of the following :

- (a) According to data released by the World Bank, the mean PM 10 (particulate matter) concentration for the city of Kabul, Afghanistan, in 1999 was 46. Suppose that because of efforts to improve air quality in Kabul, increases in modernization, and efforts to establish environment- friendly businesses,

city leaders believe rates of particulate matter in Kabul have decreased. To test this notion, they randomly sample 10 readings over a one-year period with the resulting readings shown below. Do these data present enough evidence to determine that PM 10 readings are significantly less now in Kabul? Assume that particulate readings are normally distributed and that $\alpha = .01$.

31	44	35	53	57	47	32	40	31	40
----	----	----	----	----	----	----	----	----	----


(9)

- (b) On average, 20% of the emergency room patients at Greenwood General Hospital lack health insurance. In a random sample of 4 patients, what is the probability that at least 2 will be uninsured? What is the mean and standard deviation of this binomial distribution? (9)

- (c) (i) Discuss the Properties of Mean and Standard deviation.

- (ii) Write a short note on Binomial and Poisson distributions. (9)


TABLE 1 CUMULATIVE PROBABILITIES FOR THE STANDARD NORMAL DISTRIBUTION



Entries in the table give the area under the curve to the left of the z value. For example, for $z = -.85$, the cumulative probability is .1977.

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0238	.0232	.0227	.0222	.0217	.0212	.0207	.0202	.0197	.0193
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0483	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0733	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1131	.1113	.1093	.1073	.1056	.1038	.1020	.1003	.0985	.0968
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

TABLE 1 CUMULATIVE PROBABILITIES FOR THE STANDARD NORMAL DISTRIBUTION (Continued)



Entries in the table give the area under the curve to the left of the z value. For example, for $z = 1.25$, the cumulative probability is .8944.

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990

t Table

cum. prob. one-tail	$t_{.50}$	$t_{.25}$	$t_{.20}$	$t_{.15}$	$t_{.10}$	$t_{.05}$	$t_{.025}$	$t_{.01}$	$t_{.005}$	$t_{.001}$	$t_{.0005}$
two-tail	1.00	0.50	0.40	0.30	0.20	0.10	0.05	0.02	0.01	0.002	0.001
df											
1	0.000	1.000	1.376	1.963	3.078	6.314	12.71	31.82	63.66	318.31	636.62
2	0.000	0.816	1.061	1.385	1.886	2.920	4.303	6.965	9.925	22.327	31.599
3	0.000	0.765	0.978	1.250	1.638	2.353	3.182	4.541	5.841	10.215	12.924
4	0.000	0.741	0.941	1.190	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	0.000	0.727	0.920	1.156	1.476	2.015	2.571	3.365	4.032	5.693	6.869
6	0.000	0.718	0.906	1.134	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	0.000	0.711	0.896	1.119	1.415	1.895	2.365	2.998	3.559	4.785	5.408
8	0.000	0.706	0.890	1.108	1.397	1.860	2.306	2.898	3.355	4.501	5.041
9	0.000	0.703	0.886	1.100	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	0.000	0.700	0.883	1.093	1.372	1.812	2.229	2.764	3.169	4.144	4.557
11	0.000	0.697	0.879	1.088	1.363	1.796	2.201	2.718	3.105	4.025	4.437
12	0.000	0.695	0.877	1.083	1.358	1.782	2.179	2.681	3.055	3.930	4.318
13	0.000	0.694	0.876	1.079	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	0.000	0.692	0.868	1.076	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	0.000	0.691	0.866	1.074	1.341	1.753	2.131	2.602	2.947	3.739	4.073
16	0.000	0.690	0.865	1.071	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	0.000	0.689	0.863	1.069	1.333	1.740	2.110	2.567	2.897	3.640	3.965
18	0.000	0.689	0.862	1.067	1.330	1.734	2.101	2.552	2.878	3.610	3.922
19	0.000	0.688	0.861	1.066	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	0.000	0.687	0.860	1.064	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	0.000	0.686	0.859	1.063	1.323	1.721	2.080	2.516	2.831	3.527	3.819
22	0.000	0.686	0.858	1.061	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	0.000	0.685	0.858	1.060	1.319	1.714	2.069	2.500	2.807	3.485	3.768
24	0.000	0.685	0.857	1.059	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	0.000	0.684	0.856	1.058	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	0.000	0.684	0.856	1.058	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	0.000	0.684	0.855	1.057	1.314	1.703	2.052	2.473	2.771	3.421	3.689
28	0.000	0.683	0.855	1.056	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	0.000	0.683	0.854	1.055	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	0.000	0.683	0.854	1.055	1.310	1.697	2.042	2.457	2.750	3.385	3.646
40	0.000	0.681	0.851	1.050	1.303	1.684	2.021	2.423	2.704	3.337	3.551
60	0.000	0.679	0.848	1.045	1.296	1.671	2.000	2.390	2.660	3.232	3.480
80	0.000	0.678	0.846	1.043	1.292	1.664	1.990	2.374	2.639	3.195	3.416
100	0.000	0.677	0.845	1.042	1.290	1.660	1.984	2.364	2.626	3.174	3.390
1000	0.000	0.675	0.842	1.037	1.282	1.645	1.962	2.330	2.581	3.098	3.300
Z	0.000	0.674	0.842	1.036	1.282	1.645	1.960	2.326	2.576	3.090	3.291
	0%	50%	60%	70%	80%	90%	95%	98%	99%	99.8%	99.9%
Confidence Level											

Critical Values from the t-distribution

df	t_1	$t_{.05}$	$t_{.025}$	$t_{.01}$	$t_{.005}$	$t_{.001}$
1	3.078	6.314	12.706	31.821	63.657	318.309
2	1.886	2.920	4.303	6.965	9.925	22.327
3	1.638	2.353	3.182	4.541	5.841	10.215
4	1.533	2.132	2.776	3.747	4.604	7.173

5	1.476	2.015	2.571	3.365	4.032	5.893
6	1.440	1.943	2.447	3.143	3.707	5.208
7	1.415	1.895	2.365	2.998	3.499	4.785
8	1.397	1.860	2.306	2.896	3.355	4.501
9	1.383	1.833	2.262	2.821	3.250	4.297
10	1.372	1.812	2.228	2.764	3.169	4.144
11	1.363	1.796	2.201	2.718	3.106	4.025
12	1.356	1.782	2.179	2.681	3.055	3.930
13	1.350	1.771	2.160	2.650	3.012	3.852
14	1.345	1.761	2.145	2.624	2.977	3.787
15	1.341	1.753	2.131	2.602	2.947	3.733
16	1.337	1.746	2.120	2.583	2.921	3.686
17	1.333	1.740	2.110	2.567	2.898	3.646
18	1.330	1.734	2.101	2.552	2.878	3.610
19	1.328	1.729	2.093	2.539	2.861	3.579
20	1.325	1.725	2.086	2.528	2.845	3.552
21	1.323	1.721	2.080	2.518	2.831	3.527
22	1.321	1.717	2.074	2.508	2.819	3.505
23	1.319	1.714	2.069	2.500	2.807	3.485
24	1.318	1.711	2.064	2.492	2.797	3.467
25	1.316	1.708	2.060	2.485	2.787	3.450
26	1.315	1.706	2.056	2.479	2.779	3.435
27	1.314	1.703	2.052	2.473	2.771	3.421
28	1.313	1.701	2.048	2.467	2.763	3.408
29	1.311	1.699	2.045	2.462	2.756	3.396
30	1.310	1.697	2.042	2.457	2.750	3.385
35	1.306	1.690	2.030	2.438	2.724	3.340
40	1.303	1.684	2.021	2.423	2.704	3.307
50	1.299	1.676	2.009	2.403	2.678	3.261
60	1.296	1.671	2.000	2.390	2.660	3.232
70	1.294	1.667	1.994	2.381	2.648	3.211
80	1.292	1.664	1.990	2.374	2.639	3.195
90	1.291	1.662	1.987	2.368	2.632	3.183
100	1.290	1.660	1.984	2.364	2.626	3.174
∞	1.282	1.645	1.960	2.326	2.576	3.090

Table 4 Binomial Probability Distribution $C_{nr} p^r q^{n-r}$ This table shows the probability of r successes in n independent trials, each with probability of success p .

n	r	p																			
		.01	.05	.10	.15	.20	.25	.30	.35	.40	.45	.50	.55	.60	.65	.70	.75	.80	.85	.90	.95
2	0	.980	.902	.810	.723	.640	.563	.490	.423	.360	.303	.250	.203	.160	.123	.090	.063	.040	.023	.010	.002
	1	.020	.095	.180	.255	.320	.375	.420	.455	.480	.495	.500	.495	.480	.455	.420	.375	.320	.255	.180	.095
	2	.000	.002	.010	.023	.040	.063	.090	.123	.160	.203	.250	.303	.360	.423	.490	.563	.640	.723	.810	.902
3	0	.970	.857	.729	.614	.512	.422	.343	.275	.218	.166	.125	.091	.064	.043	.027	.016	.008	.003	.001	.000
	1	.029	.135	.243	.325	.384	.422	.441	.444	.432	.408	.375	.334	.288	.239	.189	.141	.098	.057	.027	.007
	2	.000	.007	.027	.057	.096	.141	.189	.239	.288	.334	.375	.408	.432	.444	.441	.422	.384	.325	.243	.135
	3	.000	.000	.001	.003	.008	.016	.027	.043	.064	.091	.125	.166	.216	.275	.343	.422	.512	.614	.729	.857
4	0	.961	.815	.656	.522	.410	.316	.240	.179	.130	.092	.062	.041	.026	.015	.008	.004	.002	.001	.000	.000
	1	.039	.171	.292	.368	.410	.422	.412	.384	.348	.300	.250	.200	.154	.112	.076	.047	.026	.011	.004	.000
	2	.001	.014	.049	.098	.154	.211	.265	.311	.346	.368	.375	.368	.346	.311	.265	.211	.154	.098	.049	.014
	3	.000	.000	.004	.011	.026	.047	.076	.112	.154	.200	.250	.300	.346	.384	.412	.422	.410	.368	.292	.171
	4	.000	.000	.000	.001	.002	.004	.008	.015	.026	.041	.062	.092	.130	.179	.240	.316	.410	.522	.656	.815
5	0	.951	.774	.590	.444	.328	.237	.168	.116	.078	.050	.031	.019	.010	.005	.002	.001	.000	.000	.000	.000
	1	.048	.204	.328	.392	.410	.396	.360	.312	.259	.206	.156	.113	.077	.049	.028	.015	.006	.002	.000	.000
	2	.001	.021	.073	.138	.205	.264	.309	.336	.346	.337	.312	.276	.230	.181	.132	.088	.051	.024	.008	.001
	3	.000	.001	.008	.024	.051	.088	.132	.181	.230	.276	.312	.337	.346	.336	.309	.264	.205	.138	.073	.021
	4	.000	.000	.000	.002	.006	.015	.028	.049	.077	.113	.156	.206	.259	.312	.360	.396	.410	.392	.328	.204
	5	.000	.000	.000	.000	.000	.001	.002	.005	.010	.019	.031	.050	.078	.116	.168	.237	.328	.444	.590	.774
6	0	.941	.735	.531	.377	.262	.178	.118	.075	.047	.028	.016	.008	.004	.002	.001	.000	.000	.000	.000	.000
	1	.057	.232	.354	.399	.393	.356	.303	.244	.187	.136	.094	.061	.037	.020	.010	.004	.002	.000	.000	.000
	2	.001	.031	.098	.176	.246	.297	.324	.328	.311	.278	.234	.188	.138	.095	.060	.033	.015	.006	.001	.000
	3	.000	.002	.015	.042	.082	.132	.185	.236	.276	.303	.312	.303	.276	.236	.185	.132	.082	.042	.015	.002
	4	.000	.000	.001	.006	.015	.033	.060	.095	.138	.186	.234	.278	.311	.328	.324	.297	.246	.176	.098	.031
	5	.000	.000	.000	.000	.002	.004	.010	.020	.037	.061	.094	.136	.187	.244	.303	.356	.393	.399	.354	.232
	6	.000	.000	.000	.000	.000	.000	.001	.002	.004	.008	.016	.028	.047	.075	.118	.178	.262	.377	.531	.735
7	0	.932	.698	.478	.321	.210	.133	.082	.049	.028	.015	.008	.004	.002	.001	.000	.000	.000	.000	.000	.000
	1	.066	.257	.372	.396	.367	.311	.247	.185	.131	.087	.055	.032	.017	.008	.004	.001	.000	.000	.000	.000
	2	.002	.041	.124	.210	.275	.311	.318	.299	.261	.214	.164	.117	.077	.047	.025	.012	.004	.001	.000	.000
	3	.000	.004	.023	.062	.115	.173	.227	.268	.290	.292	.273	.239	.194	.144	.097	.058	.029	.011	.003	.000
	4	.000	.000	.003	.011	.029	.058	.097	.144	.194	.239	.273	.292	.290	.268	.227	.173	.115	.062	.023	.004
	5	.000	.000	.000	.001	.004	.012	.025	.047	.077	.117	.164	.214	.261	.299	.318	.311	.275	.210	.124	.041
	6	.000	.000	.000	.000	.000	.001	.004	.008	.017	.032	.055	.087	.131	.185	.247	.311	.367	.396	.372	.257
	7	.000	.000	.000	.000	.000	.000	.000	.001	.002	.004	.008	.015	.028	.049	.082	.133	.210	.321	.478	.698